


AUTHOR QUERY FORM

	Journal: OO Article Number: 2123	Please e-mail or fax your responses and any corrections to: E-mail: corrections.eseo@elsevier.sps.co.in Fax: +31 2048 52799
---	---	---

Dear Author,

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>.

Articles in Special Issues: Please ensure that the words 'this issue' are added (in the list and text) to any references to other articles in this Special Issue.

Uncited references: References that occur in the reference list but not in the text – please position each reference in the text or delete it from the list.

Missing references: References listed below were noted in the text but are missing from the reference list – please make the list complete or remove the references from the text.

Location in article	Query / remark Please insert your reply or correction at the corresponding line in the proof
	No Queries

Electronic file usage

Sometimes we are unable to process the electronic file of your article and/or artwork. If this is the case, we have proceeded by:

☐

Scanning (parts of) your article

☐

Rekeying (parts of) your article

☐

Scanning the artwork

Thank you for your assistance.



Contents lists available at ScienceDirect

Oral Oncology

journal homepage: www.elsevier.com/locate/oraloncology



Review

Resident and bone marrow-derived mesenchymal stem cells in head and neck squamous cell carcinoma

Astrid De Boeck^a, Kishan Narine^b, Wilfried De Neve^c, Marc Mareel^a, Marc Bracke^a, Olivier De Wever^{a,*}

^aLaboratory of Experimental Cancer Research, Department of Radiation Oncology and Experimental Cancer Research, Ghent University Hospital, De Pintelaan 185, B-9000 Gent, Belgium

^bDivision of Cardiac Surgery, Department of Cardiac Sciences, Libin Cardiovascular Institute of Alberta, University of Calgary, Health Sciences Centre, 3330 Hospital Dr. NW, Calgary, AL, Canada T2N 4N1

^cDepartment of Radiation Oncology, Ghent University Hospital, De Pintelaan 185, 9000 Gent, Belgium

ARTICLE INFO

Article history:

Received 29 October 2009
Received in revised form 21 January 2010
Accepted 22 January 2010
Available online xxx

Keywords:

Stromal cells
Invasion
Angiogenesis
Immunomodulation
Myofibroblast

SUMMARY

Head and neck squamous cell carcinoma (HNSCC) is a major healthcare problem worldwide affecting more than half a million patients each year. Despite considerable advances in the treatment of HNSCC, a high rate of recurrences aggravates the clinical situation and disease outcomes have only modestly improved. Recent insights show that cancer is not only a disease of the transformed epithelium but is also influenced and dependent on its stromal environment. In this review we suggest that resident and bone marrow (BM)-derived mesenchymal stem cells (MSCs) are precursors of the stroma associated with HNSCC and contribute to blood and lymphangiogenesis, modulate the immune system and produce tumor-associated myofibroblasts. In addition, the impact of radiation therapy on the stromal reaction in HNSCC is discussed. Understanding the mechanisms of how MSCs promote invasive growth and metastasis in HNSCC and respond to cancer management strategies is of profound medical importance and will help us to design improved therapeutic protocols.

© 2010 Published by Elsevier Ltd.

Introduction

Head and neck squamous cell carcinoma (HNSCC) primarily affects the mucosa of the upper aerodigestive tract, comprising the nasal and paranasal sinuses, nasopharynx, oropharynx, oral cavity, hypopharynx and larynx. The American Cancer Society estimates that approximately 35,720 new cases of HNSCC will have been diagnosed in the United States in 2009.¹ Worldwide, HNSCC is the sixth most common malignancy with an incidence of 644,000 new cases a year.² Despite considerable advances in the treatment of HNSCC, a high rate of recurrences and distant metastasis aggravates the clinical situation.³ Recent insights show that cancer is not only a disease of the transformed epithelium but is also fundamentally influenced by its stromal environment.⁴ Pre-cancerous conditions of HNSCC, such as oral submucous fibrosis, are characterized by accumulation of type I collagen within the subepithelial tissue.⁵ In agreement, molecular classification of HNSCC using patterns of gene expression reveals distinct subtypes and includes a normal epithelium-like subtype and a subtype with high levels of antioxidant enzymes both with better recurrence-free survival data compared to subtypes with an epidermal growth factor receptor (EGFR)-pathway signature or a mesenchymal-enriched subtype.⁶

High levels of EGFR expression in squamous cell carcinoma (SCC) correlate with worse clinical outcome,⁷ and decreased response to radiotherapy, as evidenced by increased locoregional recurrence.⁸ Cetuximab is an IgG1 monoclonal antibody that exclusively targets EGFR with high affinity, and inhibits endogenous ligand binding, thereby blocking receptor dimerisation, tyrosine kinase phosphorylation, and signal transduction.⁹ A recent randomized trial showed that cetuximab plus radiotherapy (versus radiotherapy alone) significantly improves locoregional control and 5-years survival without worsening radiotherapy-related toxicity.¹⁰ The mechanisms by which cancer cells manipulate their local stroma in the mesenchymal-enriched subtype is more a matter of debate compared to the EGFR-pathway subtype. In this review we discuss that resident and bone marrow (BM)-derived mesenchymal stem cells (MSCs) are precursors of the stroma associated with HNSCC. Here, MSCs contribute to blood and lymph angiogenesis, modulate the immune system and produce tumor-associated myofibroblasts. Radiation therapy is a mainstay of curative therapy for HNSCC. Recent advances have focused primarily on fractionation schedules and the use of intensity modulated radiation therapy (IMRT), a form of high-precision radiotherapy that delivers radiation more precisely to the tumor while sparing the surrounding normal tissues. IMRT has greatly improved locoregional tumor control for paranasal sinuses and pharyngolaryngeal carcinoma but had little effect on distant metastasis.¹¹ Reports indicate that radiation

* Corresponding author. Tel.: +32 9 3323008; fax: +32 9 3324991.
E-mail address: olivier.dewever@ugent.be (O. De Wever).

increases local and distant recruitment of MSCs into irradiated tissues.^{12,13} The consequent production of factors derived from MSCs or from differentiated myofibroblasts, known to possess pro-invasive activities,¹⁴ may contribute to distant metastasis and so neutralize the benefit of locoregional control. Several reports of HNSCC suggest that there are indeed certain circumstances, not yet fully understood, under which radiotherapy favours relapse and metastasis (reviewed in¹⁵). Understanding the molecular biology of cancer progression and management in this regard motivated this detailed analysis of HNSCC-associated MSCs.

Definition and characterization of MSCs

The concept of MSCs can be traced to the late nineteenth century work of E. Goujon (a.d. 1869), confirmed by A. Baikow (a.d. 1870), who described the osteogenic potential of heterotopic transplants of rabbit BM.¹⁶ This osteogenic potential of BM was a feature of a specific subgroup of cells, termed the “Colony Forming Unit-fibroblasts” (CFU-f), which made up a very small percentage of the total BM cell population.¹⁷ Subsequent studies demonstrated that these cells could differentiate into various other mesenchymal cell lineages, and they were therefore called MSCs.¹⁸ The definition and designation of MSCs remains a point of discussion, especially since our knowledge is solely based on the characterization of cultured cells. In this review, the designation “MSCs” refers to tissue culture-adherent stromal cells isolated from a variety of tissues and capable of differentiating into cell lineages of mesenchymal tissues such as adipocytes, osteocytes, chondrocytes and connective tissue cells.

MSCs express a variety of antigens that are also expressed by many other cell types and to date no unique MSC immunophenotyping marker(s) has(ve) been identified.¹⁹ MSCs express CD73 (ecto-5'-nucleotidase), CD90 (Thy-1) and CD105 (endoglin), but not CD11b, CD14, CD19, CD34, CD45, CD79 α and HLA class II.

Classically, MSCs are plated on tissue-culture substrates in low-glucose (1 g L⁻¹) Dulbecco's Modified Eagle's Medium (DMEM), supplemented with 10% of selected batches of fetal bovine serum (FBS). After several passages, MSCs enter senescence, with changes in morphology and a reduced proliferation and differentiation potential. The pace of senescence is affected by the culture conditions. A culture system which allows dynamic expansion of a high-extension silicone rubber (HESR) substrate with a much lower stiffness (10–20 $\times 10^3$ Pa) as compared to tissue-culture substrates (2.78 $\times 10^9$ Pa),²⁰ reduces contact inhibition and results in longer preservation of the cell phenotype. Moreover, the growth on expandable HESR matrices suppresses expression of α -smooth muscle actin (SMA), a functional marker for fibrogenic myofibroblasts, expressed by MSCs on tissue-culture substrates.

MSCs in noncancerous normal and pathological situations

Embryogenesis of MSCs

MSCs in the head and neck region possibly have a different origin as compared to the rest of the body. Cranial skeleton and other mesenchymal tissues of head and neck are mainly derived from the neural crest (NC),²¹ except for the occipital and otic (partly) regions of the skull, which are derived from the mesoderm.²² Trunk and limb mesenchyme is derived from the mesoderm.²³ NC can generate MSCs. Mesoderm can generate mesenchymal tissues without transiting through an MSC intermediate and it is unclear whether MSCs are derived from the mesoderm.²⁴ The development of MSCs arises in multiple waves from distinct origins (Fig. 1).²⁴ The earliest wave originates in the neuroepithelial and NC cells. An NC gene network regulates EMT (epithelial-to-mesenchymal transition) of

neuroectodermal cells at the dorsal aspect of the neural tube and generates the NC, containing highly invasive cells that give rise to MSCs, as well as neurons and glial cells.^{21,25} Later, a wave of MSCs from as yet unidentified sources, possibly mesoderm or NC-derived MSCs, becomes increasingly important.²⁴

Besides their role in embryogenesis of mesenchymal tissues, MSCs colocalize with foci of haematopoiesis early in ontogeny suggesting that they support fetal haematopoiesis. MSCs circulate in fetal blood, from at least 7 weeks gestation at the onset of haematopoiesis and disappear from the circulation by the end of the first trimester, before haematopoiesis becomes established.²⁶

MSC distribution and function in adult tissues

BM serves as a reservoir for MSCs, were they represent 0.01–0.001% of all nucleated cells.²⁷ MSCs are also distributed throughout the body. Specifically, MSCs have been isolated from several oral tissues including dental pulp, dental follicle, apical papilla, periodontal ligament and palatine tonsil.^{28–32} MSCs are not found in peripheral blood under normal conditions,^{33–35} but can be derived from granulocyte colony-stimulating factor (G-CSF) mobilised peripheral blood or umbilical cord blood.³³ The distribution of MSCs throughout the body raised the question whether there exist a common MSC niche.³⁴ The derivation from the aorta, vena cava and other vessels points to a perivascular niche.^{34,36} With the use of the markers Stro-1 and CD146, MSCs are found lining blood vessels in human BM and dental pulp.²⁹ Localisation of MSCs to perivascular niches throughout the body gives them easy access to all tissues when needed for tissue repair or remodelling, provided they conserve the invasive characteristics of their progenitors, e.g. NC cells. MSCs can either provide daughter cells that differentiate and then participate in the structural repair of a wound, or can supply secreted factors that support wound repair and modulate the immune system.³⁷ After systemic administration, BM MSCs home and engraft in damaged organs such as vascular tissue, myocardium, brain, liver, kidney, lung and skin resulting in morphological and functional improvements (reviewed in³⁸). During radiotherapy, damage will occur in normal tissues lying in the radiation field. For radiotherapeutic treatment of HNSCC, the salivary glands are one of the tissues at risk. G-CSF-mobilized BM-derived cells specifically homed to radiation-induced damaged salivary glands after radiotherapy and induced repair processes.³⁹

MSCs in HNSCC

Tumor metastasis involves extensive interactions of the invasive cancer cells with host stromal components. Tumor stroma comprises extracellular matrix (ECM) and a plethora of cells that work in concert such as myofibroblasts, tumor-associated macrophages, mast cells, neutrophils, endothelial cells, and bacteria.⁴⁰ All of them critically influence the process of carcinogenesis and tumor progression. The mechanisms by which cancer cells manipulate their local ecosystem are still a matter of debate. In this review we discuss the hypothesis that resident and BM MSCs are precursors of the stroma associated with HNSCC, thereby promoting invasive tumor growth and distant metastasis. Furthermore, we will address the question whether radiotherapy affect MSC recruitment and differentiation and by inference the surrogate endpoints invasion and metastasis. HNSCC is a significant health problem, with extremely poor outcomes and significant morbidity if patients have a disease recurrence at the locoregional site. After surgical resection, microscopic cancer cells left behind in the wound margins of the surgical resection bed increase the likelihood of local failures.⁴¹ The immediate postoperative period may be a time of maximum growth stimulus for any residual cancer




Figure 1 Early embryonic sources and fate of MSCs. Embryonic MSCs are derived from the NC and possibly from the mesoderm. Adult MSCs are possibly derived from derivatives of the mesoderm or from NC-derived MSCs. Solid arrows indicate the direction of development. Dashed arrows indicate possible directions of development.

(stem) cell.⁴² Furthermore, the process of wound repair after surgical extirpation ~~in the surgical wound~~ involves the recruitment of MSCs and consequent secretion of many factors that stimulate repair³⁷ and relapse if residual cancer (stem) cells are present. Post-operative adjuvant radiotherapy may further increase the recruitment of MSCs into the wound site causing a vicious cancer progression cycle. Traditionally, patients with positive margins receive postoperative radiotherapy and/or chemotherapy; however the prognosis for these patients remains poor. Approximately 75% of patients with positive surgical margins develop local recurrence following radiotherapy according to one study.⁴³

Recruitment of MSCs to HNSCC

Cytokines and growth factors secreted by tumors recruit resident and distant respondent cells such as MSCs.^{44,45} Intravenously injected green fluorescent protein (GFP)-labelled MSCs in the tail vein of tumor bearing mice are recruited to xenografts derived from several cancer cell lines including UMSSC1 HNSCC cells.¹² Altered expression of cytokines and growth factors plays a major role in the malignant transformation of many cancers including HNSCC.⁴⁶ Decreasing cytokine levels in serum are associated with response to therapy, while increasing levels are related to HNSCC progression and recurrence.⁴⁷ Over-expression of tumor necrosis factor (TNF)- α , vascular endothelial growth factor (VEGF), hepatocyte growth factor (HGF), platelet-derived growth factor (PDGF)-AB, transforming growth factor (TGF)- β 1 and interleukins (ILs), has been observed in HNSCC cells in vitro as well as in patients' tumor specimens and serum.^{46,48} Among them, PDGF-AB, VEGF, HGF and IL-8 exert strong chemotactic effects on BM MSCs and are possibly involved in MSC recruitment to HNSCC.⁴⁹ Priming of MSCs with pro-inflammatory cytokines like TNF α enhances migration of MSCs in vitro suggesting that the mobilisation and subsequent homing to tumors depend on the systemic and local inflammatory state.⁴⁹ Indeed, a chronic increase in inflammatory mediators in the oral cavity and oropharynx can lead to increased invasion and metastasis.⁵⁰ Furthermore, inflammatory cytokines stimulate specific matrix metalloproteinase (MMP) activity in MSCs assisting passage through the basement membrane during extravasation.⁵¹

Anti-cancer treatment influences recruitment of MSCs. Irradiated tumors, compared to unirradiated tumors, show an increase in MSC recruitment.¹³ This was demonstrated by bilateral hind leg breast tumor implants: one was left untreated, whereas the other was irradiated before intravenous injection of MSCs. At 48 h postirradiation, more MSCs were detected in the irradiated than in the unirradiated limbs. In unirradiated tumors, MSCs were more commonly associated with intravascular or perivascular structures, whereas in irradiated tumors, MSCs were present in

higher proportions in the tumor parenchyma. MSC migration to irradiated tumors may result from a dynamic interplay in which cancer cells secrete cytokines in response to radiation, leading to chemokine receptor upregulation on MSCs, and ultimately resulting in enhanced migration towards the chemokine ligand-bearing tumor. The consequent production of MSC-derived factors may contribute to relapse and metastasis. In this context it may be interesting to compare immunohistochemically, provided specific markers can be developed, the presence of MSCs and/or terminally differentiated myofibroblasts in HNSCC patients treated or not by IMRT. In a cell culture model we may understand how MSCs react to cancer management protocols including radio-chemotherapy.

MSCs and HNSCC progression

In vitro and in vivo models have shown that MSCs stimulate invasive growth of solid and haematological tumors.^{52,53} Tumor-stroma interactions are important in HNSCC pathogenesis.^{48,54,55} We suggest that MSCs are involved in HNSCC progression by: (i) supporting blood and lymph angiogenesis, (ii) modulating the immune system, and (iii) generating tumor-associated myofibroblasts (Fig. 2).

MSCs support blood and lymph angiogenesis

Blood and lymph angiogenesis are key components of the metastatic spread of cancer cells. Blood angiogenesis is controlled by angiogenic factors directly produced by cancer cells, as well as by factors from the surrounding stromal tissues. VEGF and IL-8 are prominent pro-angiogenic factors that are upregulated in HNSCC tumors and associated with aggressive tumor growth and decreased survival.⁵⁶ MSCs can act as precursors of endothelial cells and pericytes and promote angiogenesis.⁵⁷ When treated with VEGF, MSCs acquire an endothelial cell phenotype, with expression of vascular endothelial growth factor receptor (VEGFR)-1, VEGFR-2, vascular endothelial (VE)-cadherin, vascular cell adhesion molecule (VCAM)-1 and von Willebrand Factor (vWF).⁵⁸ MSCs reside in perivascular niches throughout the body^{29,34} and can engraft within blood vessels at sites of hypoxia,⁵⁹ supporting the contribution of MSCs in blood angiogenesis at hypoxic tumor sites. In addition, paracrine factors of MSCs recruit endothelial cells and smooth muscle cells and stimulate their proliferation.⁶⁰ Local injection of MSC-derived conditioned medium (CM) enhances vascularisation and perfusion in an ischemic hindlimb mouse model.⁶⁰ Tissue hypoxia, as present in tumors, is a major stimulus for vascularisation. Hypoxia stimulates MSCs to a 2-fold increase in secretion of pro-angiogenic factors like VEGF-A, fibroblast growth factor (FGF)-2, FGF-7, IL-1, IL-6, PDGF, TGF- β , TNF- α .^{60–62} Hypoxic MSCs show




Figure 2 Resident and BM-derived MSCs support invasive tumor growth. Chemoattractants divert resident perivascular and BM-derived MSCs towards the tumor. MSCs undergo differentiation into myfibroblasts, endothelial cells and pericytes. Myfibroblasts stimulate invasion of cancer cells. Thin arrows indicate cellular or tumoral transition. Thick arrows indicate displacement or invasion. Dashed arrows indicate diffusion of soluble factors. Adapted from [110].

improved angiogenic potency compared to naive MSCs when implanted in ischemic hindlimbs.⁶²

The involvement of regional lymph nodes is an important indicator of tumor aggressiveness and is a prognostic factor for HNSCC patients. Increased tumor lymph angiogenesis correlates with lymph node metastasis in HNSCC,⁶³ but the mechanisms regulating metastatic spread through the lymphatic route remain largely unexplored to date. VEGF-C and VEGF-D are implicated in tumor lymph angiogenesis and lymph node metastases.⁶⁴ A direct correlation exist between VEGF-C expression and the presence of lymph node metastases in HNSCC.^{63,65,66} In patients with tongue carcinoma, VEGF-C was associated with primary tumor size, regional lymph node metastasis, distant metastasis and prognosis. Moreover, VEGF-C expression correlated with locoregional recurrence and distant failure.⁶⁶ MSCs have been shown to play a role in lymph angiogenesis and acquire a lymphatic phenotype when exposed to VEGF-C.⁶⁷ Migratory activity of MSCs towards VEGF-C in vitro suggests that VEGF-C may recruit circulating MSCs.

The identification of soluble cancer or stromal cell-derived mediators which stimulate both blood and lymph angiogenesis can reveal targets by which to interrupt tumor angiogenesis which would, in turn, limit the growth and metastatic potential of solid cancers such as HNSCC.

MSCs modulate the immune system

HNSCC develop molecular strategies to escape efficient antitumor immune responses.^{68,69} HNSCC are infiltrated primary with T cells and dendritic cells (DC), but also with B cells, natural killer (NK) cells, macrophages and eosinophils. Impaired function of T cells and DC is observed in HNSCC.⁶⁸ MSCs exert local immunosuppressive effects, implicating that engraftment of MSCs in HNSCC creates an immunosuppressive environment. Djouad et al.⁷⁰ demonstrated that MSCs prevented the rejection of allogenic tumor cells in immunocompetent mice. MSCs infused systemically or adjacent to subcutaneously implanted B16 melanoma cells re-

sulted in enhanced tumor formation, whereas melanoma cells injected alone were eliminated by the host immune system, suggesting a facilitatory role of MSCs on allogenic tumor formation. MSCs affect the proliferation and function of immune cells including T cells, DC, NK cells, B lymphocytes and macrophages (reviewed in⁷¹). In vivo and in vitro evidence suggest that the proliferation of stimulated T cells is inhibited by MSCs without immunological restriction; similar suppressive effects being observed under autologous and allogenic conditions.⁷² This suppression of T cells affects antigen specific proliferation, expression of activation markers, cytotoxic T cell (CTL) formation and interferon (IFN)- γ and IL-4 production.^{73,74} MSCs also induce T cell anergy, an observation supported by their lack of co-stimulatory molecules (B7-1 and B7-2) and the restoration of proliferation following MSCs removal.^{73,75,76} In addition, MSCs modulate the effects of CTLs and suppress CD8 + CTLs, but not activated CD8 + CTLs cells, suggesting a possible inhibition of lymphocyte proliferation with reduced overall CTLs cytolytic response rather than inhibiting cytotoxicity itself.^{77,78} Both human and mice MSCs inhibit the proliferation of B cells stimulated by CD40L and cytokines.^{79,80} Furthermore, human MSCs inhibit the differentiation, chemotactic behaviour and antibody secretion of B cells.⁸⁰

MSCs are a source of HNSCC-associated myfibroblasts

Myfibroblasts are abundantly present in the stroma of developing tumors and drive invasive tumor growth by providing a suitable environment.^{81,82} There is ample evidence for the pro-invasive growth activity of tumor-associated myfibroblasts (recently reviewed in¹⁴), and there is no evidence to suggest that there is a different behaviour of tumor-associated myfibroblasts between HNSCC and other tumors. Myfibroblasts are large spindle-shaped cells with indented nuclei, α -SMA containing stress fibers and well-developed cell-matrix interactions (fibronexus). Characterization of stromal myfibroblasts is based on a combination of positive markers such as α -SMA, γ -SMA, desmin, vimentin, prolyl-4

Table 1

Efferent signals leading to the upregulation of myofibroblast markers and pro-invasive molecules in MSCs.

Efferent signal	Myofibroblast marker	Pro-invasive molecule	Reference
Single agent	1-Oleol-LPA	α -SMA	106
	5-Azacytidine	α -SMA; desmin; FSP	92
	PDGF-AA	α -SMA	107
	TGF- β 1	α -SMA; calponin; SM22 α	106
	TGF- β 3; D-erythro-SPC	α -SMA; calponin; SM22 α	108
CCCM	MDA-MB-231; PANC-1; U87	α -SMA; desmin; FSP	92
	OVCAR-3; ascites ovary tumor	α -SMA	106
	SK-OV-3	α -SMA; desmin; FAP; FSP	52
		FAP; IL-6; TGF- β 1; TNC; TSP1; VEGF	
Coculture	HCT115; HT29	α -SMA; calponin	109

Abbreviations: CCCM, cancer cell-conditioned medium; LPA, lysophosphatidic acid; PDGF, platelet-derived growth factor; TGF, tumor growth factor; SPC, sphingosylphosphorylcholine; SMA, smooth muscle actin; FSP, fibroblast-specific protein; SM22 α , smooth muscle 22 α ; FAP, fibroblast activating protein; SDF, stromal-derived factor; TNC, tenascin-C; TSP1, thrombospondin-1; IL, interleukin; VEGF, vascular endothelial growth factor.

hydroxylase (P4H) and negative markers such as cytokeratin, CD31, CD34 and [smoothelin](#).¹⁴ Stromal myofibroblasts produce ECM components, ECM remodelling enzymes, growth factors, cytokines and chemokines in order to create an invasive growth promoting [ecosystem](#).¹⁴

Immunohistochemical analysis of α -SMA reveals that myofibroblasts are abundantly present in the stroma of [HNSCC](#).^{54,83–85} iTRAQ (a non-gel based technique using isotopes to identify and quantify proteins from different sources in one single experiment) multidimensional liquid chromatography and tandem mass spectrometry revealed the myofibroblast-associated S100-A11 protein calgizarrin as a novel HNSCC [biomarker](#).⁸⁶ Myofibroblast appearance increases with increasing tumor invasiveness in squamous cell carcinoma of the oral [cavity](#).⁸¹ Abundant presence of myofibroblasts in the stroma is associated with several clinicopathological features of HNSCC including lymph node metastasis, disease stage and regional [recurrence](#).^{54,83} Understanding the origin and molecular events for the generation of tumor-associated myofibroblasts is still a matter of debate. Tumor-associated myofibroblasts are thought to arise from several mobilised cell types⁸⁷ including migratory neighbours such as tissue-resident MSCs or [tissue-resident](#) fibroblasts, and distant invaders such as BM-derived [MSCs](#).^{44,45,88} Myofibroblast differentiation is regulated by growth factors, mainly of the TGF- β [family](#),⁸⁹ which are secreted abundantly by [HNSCC](#).^{82,90} Interestingly, irradiation generates reactive oxygen species causing oxidation of specific amino acids in the latent TGF- β complex and release of its active [form](#).⁹¹ MSCs exposed to cancer cell-conditioned medium (CCCM), ascites from ovarian cancer patients or soluble cancer cell-derived factors like TGF- β or after long-term coculture with cancer cells, acquire a myofibroblast phenotype, characterized by an increased α -SMA expression and ECM, protease, and growth factor production ([Table 1](#)). These data put forward MSCs as myofibroblast precursors in the stroma of several solid cancers including HNSCC. Moreover, gene expression profiling reveals similarities between CCCM-exposed MSCs and stromal [myofibroblasts](#).^{92,93} The fact that myofibroblasts share surface antigens and functions with MSCs suggest they may originate from the [BM](#).⁹⁴ [BM-derived](#) MSC's contribute to 25% of the total myofibroblast population in the tumor stroma in a mouse model of pancreatic insulinoma⁴⁴ and in a subcutaneous pancreatic xenograft [tumor](#).⁴⁵ Furthermore, these [BM-derived](#) MSC-derived myofibroblasts actively participate in the production of matrix proteins, such as collagen type I, in xenograft [tumors](#).⁴⁴ TGF- β 1 derived from several oral squamous cell carcinoma cell lines (OSCC) induces myofibroblast differentiation of primary fibroblasts. Fibrosis in metastatic lymph nodes is a factor of worse prognosis in cancer of the oral [cavity](#).³⁵ Several paracrine factors produced by MSCs may be implicated in HNSCC progression. The

c-Met receptor tyrosine kinase is a potential therapeutic target for HNSCC since scatter factor (SF)/HGF, secreted by stromal HNSCC cells, stimulates invasive growth and [angiogenesis](#).^{95,96} Galectin-1 is significantly overexpressed in the tumor-associated stroma as well as in the invasion front during early oral carcinogenesis and associate with worse disease-free [survival](#).⁹⁷ Stromal-derived factor (SDF)-1 α , frequently detected in secretomes of [MSCs](#),⁹⁸ promotes invasion of HNSCC by activating Nuclear [Factor- \$\kappa\$ B](#).⁹⁹ Furthermore, gene expression profiles identify activation of Nuclear [Factor- \$\kappa\$ B](#) as characteristic of high-risk [HNSCC](#).¹⁰⁰ Activation of toll-like receptor 4 signalling in HNSCC cells promotes tumor development¹⁰¹ and this activation may be mediated by the tumor-associated stromal cell-derived ECM protein tenascin-C ([TNC](#)).¹⁰² Coculture of primary and metastatic HNSCC cells with fibroblasts derived from human gingiva causes increases in expression of cytokines which are involved in HNSCC cell [invasion](#).⁴⁸ Fibroblast-derived membrane type 1 (MT1)-MMP promote HNSCC cancer cell invasion in cell culture and tumor growth in xenograft [models](#).¹⁰³ The so called isolated fibroblast populations in these studies are likely to be multipotent MSCs, since MSCs can be isolated from a variety of oral tissues by explant culture, as used in these [studies](#).¹⁰⁴

Conclusions and perspectives

Understanding the mechanisms of how supporting host cells composing the tumor ecosystem promote invasive growth and metastasis and react to cancer management strategies is profoundly important. Given the role of MSCs in wound repair³⁷ and their emerging use as therapeutic [agents](#),¹⁰⁵ we propose that MSCs are a critical, manipulable component in pre-cancerous conditions such as oral submucous fibrosis⁵ and in the tumor ecosystem of HNSCC. Understanding the role of MSCs within pre-cancerous and cancerous conditions will be extremely valuable. What we lack are methods to specifically mark and trace the lineage of resident MSCs. Such methods, when available, will help us to determine the extent to which MSCs act as stem cells or as sources of secreted factors, as well as to identify distinct functional subpopulations.

Conflicts of Interest Statement

None [declared](#).

Acknowledgments

This work was supported by Fund for Scientific Research-Flanders (Brussels, [Belgium](#)). A. De Boeck is a doctoral researcher and O. De Wever is a post-doctoral researcher.

References

- Jemal A, Siegel R, Ward E, Hao Y, Xu J, Thun MJ. Cancer statistics, 2009. *CA Cancer J Clin* 2009;**59**(4):225–49.
- Parkin DM, Bray F, Ferlay J, Pisani P. Global cancer statistics, 2002. *CA Cancer J Clin* 2005;**55**(2):74–108.
- Haddad RI, Shin DM. Recent advances in head and neck cancer. *N Engl J Med* 2008;**359**(11):1143–54.
- Weber F, Xu YM, Zhang L, et al. Microenvironmental genomic alterations and clinicopathological behavior in head and neck squamous cell carcinoma. *JAMA* 2007;**297**(2):187–95.
- Teh MT, Tilakaratne WM, Chaplin T, et al. Fingerprinting genomic instability in oral submucous fibrosis. *J Oral Pathol Med* 2008;**37**(7):430–6.
- Chung CH, Parker JS, Karaca G, et al. Molecular classification of head and neck squamous cell carcinomas using patterns of gene expression. *Cancer Cell* 2004;**5**(5):489–500.
- He YK, Zeng Q, Drenning SD, et al. Inhibition of human squamous cell carcinoma growth in vivo by epidermal growth factor receptor antisense RNA transcribed from the U6 promoter. *J Natl Cancer Inst* 1998;**90**(14):1080–7.
- Ang KK, Berkey BA, Tu XY, et al. Impact of epidermal growth factor receptor expression on survival and pattern of relapse in patients with advanced head and neck carcinoma. *Cancer Res* 2002;**62**(24):7350–6.
- Li S, Schmitz KR, Jeffrey PD, Wiltzius JJ, Kussie P, Ferguson KM. Structural basis for inhibition of the epidermal growth factor receptor by cetuximab. *Cancer Cell* 2005;**7**(4):301–11.
- Bonner JA, Harari PM, Giralt J, et al. Radiotherapy plus cetuximab for locoregionally advanced head and neck cancer: 5-year survival data from a phase 3 randomised trial, and relation between cetuximab-induced rash and survival. *Lancet Oncol* in press;**11**(1):21–8 [[doi:10.1016/S1470-2045\(09\)70311-0](https://doi.org/10.1016/S1470-2045(09)70311-0)].
- Gregoire V, De Neve W, Eisbruch A, Lee N, Van den Weyngaert D, Van Gestel D. Intensity-modulated radiation therapy for head and neck carcinoma. *Oncologist* 2007;**12**(5):555–64.
- Zielske SP, Livant DL, Lawrence TS. Radiation increases invasion of gene-modified mesenchymal stem cells into tumors. *Int J Radiat Oncol Biol Phys* 2009;**75**(3):843–53.
- Klopp AH, Spaeth EL, Dembinski JL, et al. Tumor irradiation increases the recruitment of circulating mesenchymal stem cells into the tumor microenvironment. *Cancer Res* 2007;**67**(24):11687–95.
- De Wever O, Demetter P, Mareel M, Bracke M. Stromal myofibroblasts are drivers of invasive cancer growth. *Int J Cancer* 2008;**123**(10):2229–38.
- Madani I, De Neve W, Mareel M. Does ionizing radiation stimulate cancer invasion and metastasis? *Bull Cancer* 2008;**95**(3):292–300.
- Dennis JE, Caplan AL. Bone marrow mesenchymal stem cells. In: Sell S, editor. *Stem cells handbook*. Totowa: Humana Press; 2004. p. 107–18.
- Friedenstein AJ. Determined and inducible osteogenic precursor cells. In: Katherine Elliott DWF, editor. *Ciba foundation symposium 11 – hard tissue growth, repair and remineralization*; 1973. p. 169–85.
- Caplan AL. Mesenchymal stem cells. *J Orthop Res* 1991;**9**(5):641–50.
- Dominici M, Blanc KL, Mueller I, et al. Minimal criteria for defining multipotent mesenchymal stromal cells. The international society for cellular therapy position statement. *Cytotherapy* 2006;**8**(4):315–7.
- Majd H, Wipff P-J, Buscemi L, et al. A novel method of dynamic culture surface expansion improves mesenchymal stem cell proliferation and phenotype. *Stem Cells* 2009;**27**(1):200–9.
- Le Douarin NM, Creuzet S, Couly G, Dupin E. Neural crest cell plasticity and its limits. *Development* 2004;**131**(19):4637–50.
- Couly GF, Coltey PM, Le Douarin NM. The triple origin of skull in higher vertebrates: a study in quail-chick chimeras. *Development* 1993;**117**(2):409–29.
- Young HE, Black Jr AC. Adult stem cells. *Anat Rec A Discov Mol Cell Evol Biol* 2004;**276**(1):75–102.
- Takashima Y, Era T, Nakao K, et al. Neuroepithelial cells supply an initial transient wave of MSC differentiation. *Cell* 2007;**129**(7):1377–88.
- Nikitina N, Sauka-Spengler T, Bronner-Fraser M. Dissecting early regulatory relationships in the lamprey neural crest gene network. *Proc Natl Acad Sci USA* 2008;**105**(51):20083–8.
- Mendes SC, Robin C, Dzierzak E. Mesenchymal progenitor cells localize within hematopoietic sites throughout ontogeny. *Development* 2005;**132**(5):1127–36.
- Pittenger MF, Mackay AM, Beck SC, et al. Multilineage potential of adult human mesenchymal stem cells. *Science* 1999;**284**(5411):143–7.
- Lindroos B, Mäenpää K, Ylikomi T, Oja H, Suuronen R, Miettinen S. Characterisation of human dental stem cells and buccal mucosa fibroblasts. *Biochem Biophys Res Commun* 2008;**368**(2):329–35.
- Shi S, Gronthos S. Perivascular niche of postnatal mesenchymal stem cells in human bone marrow and dental pulp. *J Bone Miner Res* 2003;**18**(4):696–704.
- Pierdomenico L, Bonsi L, Calvitti M, et al. Multipotent mesenchymal stem cells with immunosuppressive activity can be easily isolated from dental pulp. *Transplantation* 2005;**80**(6):836–42.
- Janjanin S, Djouad F, Shanti RM, et al. Human palatine tonsil: a new potential tissue source of multipotent mesenchymal progenitor cells. *Arthritis Res Ther* 2008;**10**(4):R83.
- Sonoyama W, Liu Y, Yamaza T, et al. Characterization of the apical papilla and its residing stem cells from human immature permanent teeth: a pilot study. *J Endod* 2008;**34**(2):166–71.
- Tondreau T, Meuleman N, Delforge A, et al. Mesenchymal stem cells derived from CD133-positive cells in mobilized peripheral blood and cord blood: proliferation, Oct4 expression, and plasticity. *Stem Cells* 2005;**23**(8):1105–12.
- Meirelles LDS, Chagastelles PC, Nardi NB. Mesenchymal stem cells reside in virtually all post-natal organs and tissues. *J Cell Sci* 2006;**119**(11):2204–13.
- Lehn CN, Rapoport A. The desmoplastic lymph node reaction as a prognostic factor of cancer of the tongue and floor of the mouth. *Sao Paulo Med J* 1994;**112**(3):591–6.
- Crisan M, Yap S, Casteilla L, et al. A perivascular origin for mesenchymal stem cells in multiple human organs. *Cell Stem Cell* 2008;**3**(3):301–13.
- Stappenbeck TS, Miyoshi H. The role of stromal stem cells in tissue regeneration and wound repair. *Science* 2009;**324**(5935):1666–9.
- Chamberlain G, Fox J, Ashton B, Middleton J. Concise review: mesenchymal stem cells: their phenotype, differentiation capacity, immunological features, and potential for homing. *Stem cells* 2007;**25**(11):2739–49.
- Lombaert IMA, Wieringa PK, Kok T, Kampinga HH, deHaan G, Coppes RP. Mobilization of bone marrow stem cells by granulocyte colony-stimulating factor ameliorates radiation-induced damage to salivary glands. *Clin Cancer Res* 2006;**12**(6):1804–12.
- Joyce JA, Pollard JW. Microenvironmental regulation of metastasis. *Nat Rev Cancer* 2009;**9**(4):239–52.
- Brennan JA, Mao L, Hruban RH, et al. Molecular assessment of histopathological staging in squamous-cell carcinoma of the head and neck. *N Engl J Med* 1995;**332**(7):429–35.
- Ragaz J. Comments on kinetics and biology of the residual cancer, and on relevant therapeutic strategies based on these phenomena. *Prog Clin Biol Res* 1990;**354B**:117–39.
- Jones AS, Hanafi ZB, Nadapalan V, Roland NJ, Kinsella A, Helliwell TR. Do positive resection margins after ablative surgery for head and neck cancer adversely affect prognosis? A study of 352 patients with recurrent carcinoma following radiotherapy treated by salvage surgery. *Br J Cancer* 1996;**74**(1):128–32.
- Direkze NC, Hodivala-Dilke K, Jeffery R, et al. Bone marrow contribution to tumor-associated myofibroblasts and fibroblasts. *Cancer Res* 2004;**64**(23):8492–5.
- Ishii G, Sangai T, Oda T, et al. Bone-marrow-derived myofibroblasts contribute to the cancer-induced stromal reaction. *Biochem Biophys Res Commun* 2003;**309**(1):232–40.
- Pries R, Nitsch S, Wollenberg B. Role of cytokines in head and neck squamous cell carcinoma. *Expert Rev Anticancer Ther* 2006;**6**(9):1195–203.
- Allen C, Duffy S, Teknos T, et al. Nuclear factor-κB-related serum factors as longitudinal biomarkers of response and survival in advanced oropharyngeal carcinoma. *Clin Cancer Res* 2007;**13**(11):3182–90.
- Koontongkaew S, Amornphimoltham P, Yapong B. Tumor-stroma interactions influence cytokine expression and matrix metalloproteinase activities in paired primary and metastatic head and neck cancer cells. *Cell Biol Int* 2009;**33**(2):165–73.
- Ponte AL, Marais E, Gally N, et al. The in vitro migration capacity of human bone marrow mesenchymal stem cells: comparison of chemokine and growth factor chemotactic activities. *Stem cells* 2007;**25**(7):1737–45.
- Wang F, Arun P, Friedman J, Chen Z, Van Waes C. Current and potential inflammation targeted therapies in head and neck cancer. *Curr Opin Pharmacol* 2009;**9**(4):389–95.
- Ries C, Egea V, Karow M, Kolb H, Jochum M, Neth P. MMP-2, MTI-MMP, and TMP-2 are essential for the invasive capacity of human mesenchymal stem cells: differential regulation by inflammatory cytokines. *Blood* 2007;**109**(9):4055–63.
- Spaeth EL, Dembinski JL, Sasser AK, et al. Mesenchymal stem cell transition to tumor-associated fibroblasts contributes to fibrovascular network expansion and tumor progression. *PLoS One* 2009;**4**(4):e4992.
- Karnoub AE, Dash AB, Vo AP, et al. Mesenchymal stem cells within tumour stroma promote breast cancer metastasis. *Nature* 2007;**449**(7162):557–63.
- Kellermann MG, Sobral LM, da Silva SD, et al. Mutual paracrine effects of oral squamous cell carcinoma cells and normal oral fibroblasts: induction of fibroblast to myofibroblast transdifferentiation and modulation of tumor cell proliferation. *Oral Oncol* 2008;**44**(5):509–17.
- Nielsen JD, Moeslund M, Wandall HH, Dabelsteen S. Influences of tumor stroma on the malignant phenotype. *J Oral Pathol Med* 2008;**37**(7):412–6.
- Eisma RJ, Spiro JD, Kreutzer DL. Role of angiogenic factors: coexpression of interleukin-8 and vascular endothelial growth factor in patients with head and neck squamous carcinoma. *Laryngoscope* 1999;**109**(5):687–93.
- Lozito TP, Kuo CK, Taboas JM, Tuan RS. Human mesenchymal stem cells express vascular cell phenotypes upon interaction with endothelial cell matrix. *J Cell Biochem* 2009;**107**(4):714–22.
- Oswald J, Boxberger S, Jørgensen B, et al. Mesenchymal stem cells can be differentiated into endothelial cells in vitro. *Stem cells* 2004;**22**(3):377–84.
- Hamou C, Callaghan MJ, Thangarajah H, et al. Mesenchymal stem cells can participate in ischemic neovascularization. *Plast Reconstr Surg* 2009;**123**(2):455–555.
- Kinnaird T, Stabile E, Burnett MS, et al. Marrow-derived stromal cells express genes encoding a broad spectrum of arteriogenic cytokines and promote in vitro and in vivo arteriogenesis through paracrine mechanisms. *Circ Res* 2004;**94**(5):678–85.
- Potier E, Ferreira E, Andriamanalijaona R, et al. Hypoxia affects mesenchymal stromal cell osteogenic differentiation and angiogenic factor expression. *Bone* 2007;**40**(4):1078–87.

62. Li T-S, Hamano K, Suzuki K, Ito H, Zempo N, Matsuzaki M. Improved angiogenic potency by implantation of ex vivo hypoxia prestimulated bone marrow cells in rats. *Am J Physiol Heart Circ Physiol* 2002;**283**(2):H468–73.
63. Beasley NJP, Prevo R, Banerji S, et al. Intratumoral lymphangiogenesis and lymph node metastasis in head and neck cancer. *Cancer Res* 2002;**62**(5):1315–20.
64. Stacker SA, Achen MG, Jussila L, Baldwin ME, Alitalo K. Lymphangiogenesis and cancer metastasis. *Nat Rev Cancer* 2002;**2**(8):573–83.
65. Kishimoto K, Sasaki A, Yoshihama Y, Mese H, Tsukamoto G, Matsumura T. Expression of vascular endothelial growth factor-C predicts regional lymph node metastasis in early oral squamous cell carcinoma. *Oral Oncol* 2003;**39**(4):391–6.
66. Tanigaki Y, Nagashima Y, Kitamura Y, Matsuda H, Mikami Y, Tsukuda M. The expression of vascular endothelial growth factor-A and -C, and receptors 1 and 3: correlation with lymph node metastasis and prognosis in tongue squamous cell carcinoma. *Int J Mol Med* 2004;**14**(3):389–95.
67. Conrad C, Niess H, Huss R, et al. Multipotent mesenchymal stem cells acquire a lymphendothelial phenotype and enhance lymphatic regeneration in vivo. *Circulation* 2009;**119**(2):281–9.
68. Whiteside TL. Immunobiology of head and neck cancer. *Cancer Metastasis Rev* 2003;**24**(1):95–105.
69. Jewett A, Head C, Cacalano NA. Emerging mechanisms of immunosuppression in oral cancers. *J Dent Res* 2006;**85**(12):1061–73.
70. Djouad F, Plence P, Bony C, et al. Immunosuppressive effect of mesenchymal stem cells favors tumor growth in allogeneic animals. *Blood* 2003;**102**(10):3837–44.
71. Rasmussen I. Immune modulation by mesenchymal stem cells. *Exp Cell Res* 2006;**312**(12):2169–79.
72. Le Blanc K, Tammik L, Sundberg B, Haynesworth SE, Ringden O. Mesenchymal stem cells inhibit and stimulate mixed lymphocyte cultures and mitogenic responses independently of the major histocompatibility complex. *Scand J Immunol* 2003;**57**(1):11–20.
73. Krampera M, Glennie S, Dyson J, et al. Bone marrow mesenchymal stem cells inhibit the response of naive and memory antigen-specific T cells to their cognate peptide. *Blood* 2003;**101**(9):3722–9.
74. Aggarwal S, Pittenger MF. Human mesenchymal stem cells modulate allogeneic immune cell responses. *Blood* 2005;**105**(4):1815–22.
75. Di Nicola M, Carlo-Stella C, Magni M, et al. Human bone marrow stromal cells suppress T-lymphocyte proliferation induced by cellular or nonspecific mitogenic stimuli. *Blood* 2002;**99**(10):3838–43.
76. Glennie S, Soeiro I, Dyson PJ, Lam EW-F, Dazzi F. Bone marrow mesenchymal stem cells induce division arrest anergy of activated T cells. *Blood* 2005;**105**(7):2821–7.
77. Rasmussen I, Ringden O, Sundberg B, Le Blanc K. Mesenchymal stem cells inhibit the formation of cytotoxic T lymphocytes, but not activated cytotoxic T lymphocytes or natural killer cells. *Transplantation* 2003;**76**(8):1208–13.
78. Karlsson H, Samarasinghe S, Ball LM, et al. Mesenchymal stem cells exert differential effects on alloantigen and virus-specific T-cell responses. *Blood* 2008;**112**(3):532–41.
79. Augello A, Tasso R, Negrini Simone M, et al. Bone marrow mesenchymal progenitor cells inhibit lymphocyte proliferation by activation of the programmed death 1 pathway. *Eur J Immunol* 2005;**35**(5):1482–90.
80. Corcione A, Benvenuto F, Ferretti E, et al. Human mesenchymal stem cells modulate B-cell functions. *Blood* 2006;**107**(1):367–72.
81. Kawashiri S, Tanaka A, Noguchi N, et al. Significance of stromal desmoplasia and myofibroblast appearance at the invasive front in squamous cell carcinoma of the oral cavity. *Head Neck* 2009;**31**(10):1346–53.
82. Lewis MP, Lygoe KA, Nystrom ML, et al. Tumour-derived TGF-beta1 modulates myofibroblast differentiation and promotes HGF/SF-dependent invasion of squamous carcinoma cells. *Br J Cancer* 2004;**90**(4):822–32.
83. Vered M, Dobriyan A, Dayan D, et al. Tumor-host histopathologic variables, stromal myofibroblasts and risk score, are significantly associated with recurrent disease in tongue cancer. *Cancer Sci in press*; **101**(1):274–80 [doi:10.1111/j.1349-7006.2009.01357].
84. Kojic N, Zidar N, Vodopivec B, Gale N. Expression of CD34, [alpha]-smooth muscle actin, and transforming growth factor [beta]1 in squamous intraepithelial lesions and squamous cell carcinoma of the larynx and hypopharynx. *Hum Pathol* 2005;**36**(1):16–21.
85. Barth PJ, Schenck zu Schweinsberg T, Ramaswamy A, Moll R. CD34+ fibrocytes, alpha-smooth muscle antigen-positive myofibroblasts, and CD117 expression in the stroma of invasive squamous cell carcinomas of the oral cavity, pharynx, and larynx. *Virchows Arch* 2004;**444**(3):231–4.
86. Ralhan R, Desouza LV, Matta A, et al. Discovery and verification of head-and-neck cancer biomarkers by differential protein expression analysis using iTRAQ labeling, multidimensional liquid chromatography, and tandem mass spectrometry. *Mol Cell Proteomics* 2008;**7**(6):1162–73.
87. De Wever O, Pauwels P, De Craene B, et al. Molecular and pathological signatures of epithelial-mesenchymal transitions at the cancer invasion front. *Histochem Cell Biol* 2008;**130**(3):481–94.
88. Muehlberg FL, Song YH, Krohn A, et al. Tissue-resident stem cells promote breast cancer growth and metastasis. *Carcinogenesis* 2009;**30**(4):589–97.
89. Narine K, De Wever O, Van Valckenborgh D, et al. Growth factor modulation of fibroblast proliferation, differentiation, and invasion: implications for tissue valve engineering. *Tissue Eng* 2006;**12**(10):2707–16.
90. Liss C, Fekete MJ, Hasina R, Lam CD, Lingen MW. Paracrine angiogenic loop between head-and-neck squamous-cell carcinomas and macrophages. *Int J Cancer* 2001;**93**(6):781–5.
91. Barcellos-Hoff M, Dix T. Redox-mediated activation of latent transforming growth factor-beta 1. *Mol Endocrinol* 1996;**10**(9):1077–83.
92. Mishra PJ, Mishra PJ, Humeniuk R, et al. Carcinoma-associated fibroblast-like differentiation of human mesenchymal stem cells. *Cancer Res* 2008;**68**(11):4331–9.
93. Menon LG, Picinich S, Koneru R, et al. Differential gene expression associated with migration of mesenchymal stem cells to conditioned medium from tumor cells or bone marrow cells. *Stem Cells* 2007;**25**(2):520–8.
94. Haniffa MA, Collin MP, Buckley CD, Dazzi F. Mesenchymal stem cells: the fibroblasts' new clothes? *Haematologica* 2009;**94**(2):258–63.
95. Knowles LM, Stabile LP, Egloff AM, et al. HGF and c-Met participate in paracrine tumorigenic pathways in head and neck squamous cell cancer. *Clin Cancer Res* 2009;**15**(11):3740–50.
96. Seiwert TY, Jagadeeswaran R, Faoro L, et al. The MET receptor tyrosine kinase is a potential novel therapeutic target for head and neck squamous cell carcinoma. *Cancer Res* 2009;**69**(7):3021–31.
97. Chiang W-F, Liu S-Y, Fang L-Y, et al. Overexpression of galectin-1 at the tumor invasion front is associated with poor prognosis in early-stage oral squamous cell carcinoma. *Oral Oncol* 2008;**44**(4):325–34.
98. Hwang JH, Shim SS, Seok OS, et al. Comparison of cytokine expression in mesenchymal stem cells from human placenta, cord blood, and bone marrow. *J Korean Med Sci* 2009;**24**(4):547–54.
99. Rehman AO, Wang CY. SDF-1alpha promotes invasion of head and neck squamous cell carcinoma by activating NF-kappaB. *J Biol Chem* 2008;**283**(29):19888–94.
100. Chung CH, Parker JS, Fang L-Y, et al. Gene expression profiles identify epithelial-to-mesenchymal transition and activation of nuclear factor-(kappa)B signaling as characteristics of a high-risk head and neck squamous cell carcinoma. *Cancer Res* 2006;**66**(16):8210–8.
101. Szczepanski MJ, Czyskowska M, Szajnik M, et al. Triggering of toll-like receptor 4 expressed on human head and neck squamous cell carcinoma promotes tumor development and protects the tumor from immune attack. *Cancer Res* 2009;**69**(7):3105–13.
102. Midwood K, Sacre S, Piccinini AM, et al. Tenascin-C is an endogenous activator of toll-like receptor 4 that is essential for maintaining inflammation in arthritic joint disease. *Nat Med* 2009;**15**(7):774–80.
103. Zhang W, Matrisian LM, Holmbeck K, Vick CC, Rosenthal EL. Fibroblast-derived MT1-MMP promotes tumor progression in vitro and in vivo. *BMC Cancer* 2006;**6**:52.
104. Sudo K, Kanno M, Mihara K, et al. Mesenchymal progenitors able to differentiate into osteogenic, chondrogenic, and/or adipogenic cells in vitro are present in most primary fibroblast-like cell populations. *Stem Cells* 2007;**25**(7):1610–7.
105. Studeny M, Marini FC, Dembinski JL, et al. Mesenchymal stem cells: potential precursors for tumor stroma and targeted-delivery vehicles for anticancer agents. *J Natl Cancer Inst* 2004;**96**(21):1593–603.
106. Jeon ES, Moon HJ, Lee MJ, et al. Cancer-derived lysophosphatidic acid stimulates differentiation of human mesenchymal stem cells to myofibroblast-like cells. *Stem cells* 2008;**26**(3):789–97.
107. Ball SG, Shuttleworth CA, Kielty CM. Platelet-derived growth factor receptor-alpha is a key determinant of smooth muscle alpha-actin filaments in bone marrow-derived mesenchymal stem cells. *Int J Biochem Cell Biol* 2007;**39**(2):379–91.
108. Jeon ES, Moon HJ, Lee MJ, et al. Sphingosylphosphorylcholine induces differentiation of human mesenchymal stem cells into smooth-muscle-like cells through a TGF-(beta)-dependent mechanism. *J Cell Sci* 2006;**119**(23):4994–5005.
109. Emura M, Ochiai A, Horino M, Arndt W, Kamino K, Hirohashi S. Development of myofibroblasts from human bone marrow mesenchymal stem cells cocultured with human colon carcinoma cells and TGF beta 1. *In Vitro Cell Dev Biol Anim* 2000;**36**(2):77–80.
110. Mareel M, Oliveira MJ, Madani I. Cancer invasion and metastasis: interacting ecosystems. *Virchows Arch* 2009;**454**(6):599–622.